

# Mathematical Tripos Part IB: Easter 1999

## Numerical Analysis – Exercise Sheet 1<sup>1</sup>

1. Calculate *all* LU factorizations of the matrix

$$A = \begin{bmatrix} 10 & 6 & -2 & 1 \\ 10 & 10 & -5 & 0 \\ -2 & 2 & -2 & 1 \\ 1 & 3 & -2 & 3 \end{bmatrix},$$

where all diagonal elements of  $L$  are one. By using one of these factorizations, find *all* solutions of the equation  $A\mathbf{x} = \mathbf{b}$  where  $\mathbf{b}^T = [-2 \ 0 \ 2 \ 1]$ .

2. Let  $A$  be a real  $n \times n$  matrix that has the factorization  $A = LU$ , where  $L$  is lower triangular with ones on its diagonal and  $U$  is upper triangular. Prove that, for every integer  $k \in \{1, 2, \dots, n\}$ , the first  $k$  rows of  $U$  span the same space as the first  $k$  rows of  $A$ . Prove also that the first  $k$  columns of  $A$  are in the  $k$ -dimensional subspace that is spanned by the first  $k$  columns of  $L$ . Hence deduce that no LU factorization of the given form exists if we have  $\text{rank } H_k < \text{rank } B_k$ , where  $H_k$  is the leading  $k \times k$  submatrix of  $A$  and where  $B_k$  is the  $n \times k$  matrix whose columns are the first  $k$  columns of  $A$ .

3. By using column pivoting if necessary to exchange rows of  $A$ , an LU factorization of a real  $n \times n$  matrix  $A$  is calculated, where  $L$  has ones on its diagonal, and where the moduli of the off-diagonal elements of  $L$  do not exceed one. Let  $\alpha$  be the largest of the moduli of the elements of  $A$ . Prove by induction on  $i$  that elements of  $U$  satisfy the condition  $|U_{i,j}| \leq 2^{i-1}\alpha$ . Then construct  $2 \times 2$  and  $3 \times 3$  nonzero matrices  $A$  that yield  $|U_{2,2}| = 2\alpha$  and  $|U_{3,3}| = 4\alpha$  respectively.

4. Calculate the Cholesky factorization of the matrix

$$\begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 1 & 3 & 1 & 0 & 0 \\ 0 & 0 & 1 & 4 & 1 & 0 \\ 0 & 0 & 0 & 1 & 5 & 1 \\ 0 & 0 & 0 & 0 & 1 & \lambda \end{bmatrix}.$$

Deduce from the factorization the value of  $\lambda$  that makes the matrix singular. Also find this value of  $\lambda$  by seeking the vector in the null-space of the matrix whose first component is one.

---

<sup>1</sup>Corrections and suggestions to these notes should be emailed to [A.Iserles@damp.cam.ac.uk](mailto:A.Iserles@damp.cam.ac.uk). All handouts are available on the WWW at the URL <http://www.damp.cam.ac.uk/user/na/PartIB/Handouts.html>.

5. Let  $A$  be an  $n \times n$  nonsingular band matrix that satisfies the condition  $A_{i,j} = 0$  if  $|i - j| > r$ , where  $r$  is small, and let Gaussian elimination *with column pivoting* be used to solve  $A\mathbf{x} = \mathbf{b}$ . Identify all the coefficients of the intermediate equations that can become nonzero. Hence deduce that the total number of additions and multiplications of the complete calculation can be bounded by a constant multiple of  $nr^2$ .

6. The iteration  $\mathbf{x}_{k+1} = H\mathbf{x}_k + \mathbf{b}$  is applied for  $k = 0, 1, \dots$ , where  $H$  is the real  $2 \times 2$  matrix

$$H = \begin{bmatrix} \alpha & \gamma \\ 0 & \beta \end{bmatrix},$$

with  $\gamma$  large and  $|\alpha| < 1$ ,  $|\beta| < 1$ . Calculate the elements of  $H^k$  and show that they tend to zero as  $k \rightarrow \infty$ . Further, establish the equation  $\mathbf{x}_k - \mathbf{x}^* = H^k(\mathbf{x}_0 - \mathbf{x}^*)$ , where  $\mathbf{x}^*$  is defined by  $\mathbf{x}^* = H\mathbf{x}^* + \mathbf{b}$ . Thus deduce that the sequence  $\{\mathbf{x}_k\}_{k=0}^{\infty}$  converges to  $\mathbf{x}^*$ .

7. For some choice of  $\mathbf{x}_0$  the iterative method

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{x}_{k+1} + \begin{bmatrix} 0 & 0 & 0 \\ \xi & 0 & 0 \\ \eta & \zeta & 0 \end{bmatrix} \mathbf{x}_k = \mathbf{b}$$

is applied for  $k = 0, 1, \dots$ , in order to solve the linear system

$$\begin{bmatrix} 1 & 1 & 1 \\ \xi & 1 & 1 \\ \eta & \zeta & 1 \end{bmatrix} \mathbf{x} = \mathbf{b},$$

where  $\xi$ ,  $\eta$  and  $\zeta$  are constants. Find all values of the constants such that the sequence  $\{\mathbf{x}_k\}_{k=0}^{\infty}$  converges for every  $\mathbf{x}_0$  and  $\mathbf{b}$ . Give an example of nonconvergence when  $\xi = \eta = \zeta = -1$ . Is the solution always found in at most two iterations when  $\xi = \zeta = 0$ ?

8. Let  $\mathbf{a}_1$ ,  $\mathbf{a}_2$  and  $\mathbf{a}_3$  denote the columns of the matrix

$$A = \begin{bmatrix} 6 & 6 & 1 \\ 3 & 6 & 1 \\ 2 & 1 & 1 \end{bmatrix}.$$

Apply the Gram–Schmidt procedure to  $A$ , which generates orthonormal vectors  $\mathbf{q}_1$ ,  $\mathbf{q}_2$  and  $\mathbf{q}_3$ . Note that this calculation provides real numbers  $R_{k,\ell}$  such that  $\mathbf{a}_k = \sum_{\ell=1}^k R_{k,\ell} \mathbf{q}_\ell$ ,  $k = 1, 2, 3$ . Hence express  $A$  as the product  $A = QR$ , where  $Q$  and  $R$  are orthogonal and upper-triangular matrices respectively.

9. Calculate the QR factorization of the matrix of Exercise 8 by using three Givens rotations. Explain why the initial rotation can be any one of the three types

$\Omega^{(1,2)}$ ,  $\Omega^{(1,3)}$  and  $\Omega^{(2,3)}$ . Prove that the final factorization is independent of this initial choice in exact arithmetic, provided that we satisfy the condition that in each row of  $R$  the leading nonzero element is positive.

10. Let  $A$  be an  $n \times n$  matrix, and for  $i = 1, 2, \dots, n$  let  $k(i)$  be the number of zero elements in the  $i$ th row of  $A$  that come before all nonzero elements in this row and before the diagonal element  $A_{i,i}$ . Show that the QR factorization of  $A$  can be calculated by using at most  $\frac{1}{2}n(n-1) - \sum k(i)$  Givens rotations. Hence show that, if  $A$  is an upper triangular matrix except that there are nonzero elements in its first column, i.e.  $A_{i,j} = 0$  when  $2 \leq j < i \leq n$ , then its QR factorization can be calculated by using only  $2n - 3$  Givens rotations.

11. Calculate the QR factorization of the matrix of Exercise 8 by using two Householder reflections. Show that, if this technique is used to generate the QR factorization of a general  $n \times n$  matrix  $A$ , then the computation can be organised so that the total number of additions and multiplications is bounded above by a constant multiple of  $n^3$ .

12. Let

$$A = \begin{bmatrix} 3 & 4 & 7 & -2 \\ 5 & 4 & 9 & 3 \\ 1 & -1 & 0 & 3 \\ 1 & -1 & 0 & 0 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 11 \\ 29 \\ 16 \\ 10 \end{bmatrix}.$$

Calculate the QR factorization of  $A$  by using Householder reflections. In this case  $A$  is singular and you should choose  $Q$  so that the last row of  $R$  is zero. Hence identify all the least squares solutions of the inconsistent system  $A\mathbf{x} = \mathbf{b}$ , where we require  $\mathbf{x}$  to minimize  $\|A\mathbf{x} - \mathbf{b}\|_2$ . Verify that all the solutions give the same vector of residuals  $A\mathbf{x} - \mathbf{b}$ , and that this vector is orthogonal to the columns of  $A$ . There is no need to calculate the elements of  $Q$  explicitly.